

Zinc 750

Zn-based flow

Composition

- ▶ 35 % total zinc (Zn), that is 600 g/l

Specifications

- ▶ Product pH: 8 – 9
- ▶ pH at 0.5 % dilution: 8.5 ± 0.5
- ▶ Store: at +5°C minimum
- ▶ Density: 1.7

Packaging

- ▶ 1 l can, 12 l box
- ▶ 5 l can, 20 l box
- ▶ 10 l canister

Use

- ▶ Foliar application

EC FERTILISER. Zinc hydroxide solution. Zinc 750 is particularly recommended for crops with very high zinc requirements. Early application improves product effectiveness.

Dosage

- ▶ Corn: 1 x 3 l/ha, by direct soil application, 1.5 l/ha by foliar application, at 6-leaf stage
- ▶ Vegetables: 1.5 l/ha when plants reach 5 cm, repeat a fortnight later if needed
- ▶ Citrus and orchards: 100 cc/hl, repeat 2 to 3 times, or 500 cc/hl when leaves fall
- ▶ Cereals: 1.5 l/ha, at tillering stage
- ▶ Grassland: 5 l/ha, apply during growth period. Wait 10 days before grazing. To be repeated if needed.
- ▶ Rice: soak in a 1 % solution before transplanting
- ▶ To be used only where there is a recognized need. Do not exceed the appropriate dose rates.

